

- Seven-day crisis stabilization, residential care, 15 maximum census
- Male/female ages 10-17
- Gap Level of Care: Higher than outpatient, not as intense as inpatient
- Diversity of criteria/referral concerns, primarily related to disruption of functioning associated with suicidality, defiance in home/school, dangerous behavior (run away, self-injurious behavior, online promiscuity, etc.)
- Services at no cost to client/guardian
- Initial phone or walk-in crisis screening & assessment by master's level clinician
- Group therapy (2 hours/day)
- Individual therapy with master's level clinician (at least 1x/week)
- Physician assessment & possible medication review/adjustment
- Tristesse Grief Center and HIV/STD education groups (1x/weekly)
- Tulsa Public Schools teacher on-site (3 hours/school day)
- Strong decrease in depression & anxiety symptoms in one week
 - o CES-DC (depression): 45% reduction
 - o GAD-7(anxiety): 53% reduction

Three Major Goals:

A pause: Provide children a safe environment to be less impacted by community or family stressors, to be exposed to and potentially integrate new coping skills, and to talk to experienced staff;

A perspective: Clinical feedback and perceptions from experienced team members, physicians, nurses, and therapists regarding client's mental health, diagnoses, and future needs:

A plan: Assistance in connecting the parent/guardian with the next step in treatment.